

# FOOT & ANKLE INJURIES IN YOUNG ATHLETES

## WHAT PARENTS NEED TO KNOW

**10%** of all injuries seen in the ER are ankle sprains


**83%** of ankle injuries are diagnosed as ligament sprains with incomplete tears

**90%** **Not all fields are created equal:** of ankle sprains are due to poor playing field conditions

Flat feet or a rigid arch and the use of improper or worn-out footwear may **increase the risk of developing shin splints**

In football, the foot and ankle account for **16%** of all injuries

Overuse injuries include **Achilles tendonitis and stress fractures**



Rubber cleats are not necessary for **children under 10** – use an all-purpose sports shoe

Metal spikes should not be used until **teenage years**

The ankle is the **second most common site of injury in baseball players** after the shoulder

Pediatric heel pain is very different from adult heel pain; **it does not get better with walking around**

Generally, injuries seen in younger athletes fall in three categories:

**1** Injuries related to growth

**2** Overuse injuries

**3** Acute presentations

## When is a sprain really bad?

Sprains are classified in three categories:

FIRST DEGREE	SECOND DEGREE	THIRD DEGREE
<p><b>Tissue is only stretched</b></p> <ul style="list-style-type: none"> <li>Slight swelling</li> <li>Mild loss of range of motion and strength</li> <li>No decrease in stability</li> </ul>	<p><b>Involves stretching and some tearing of tissue</b></p> <ul style="list-style-type: none"> <li>Moderate swelling</li> <li>Usually includes some bruising</li> <li>Moderate loss of range of motion and strength</li> <li>Some decrease in stability</li> </ul>	<p><b>Complete tearing of tissue</b></p> <ul style="list-style-type: none"> <li>Significant swelling and bruising</li> <li>Near complete loss of range of motion and strength</li> <li>Marked decrease in stability</li> </ul>

## The heel is where it's at

**Sever's disease is a common cause of heel pain in growing kids, physically active kids.**



The most important thing to know about Sever's disease is that, with proper care, the condition usually goes away in under two months. Early diagnosis, proper care, and taking measures to protect the heel can help protect against future problems.

For more health information and tips, visit [FootHealthFacts.org](http://FootHealthFacts.org)—the patient education website of the American College of Foot and Ankle Surgeons.



**American College of Foot and Ankle Surgeons®**

*Proven leaders. Lifelong learners. Changing lives.*